

MIT BAG IN NON-SMOOTH CONVEX DOMAINS

KONSTANTIN PANKRASHKIN

Dedicated to the memory of my doctoral co-advisor Jochen Brüning (1947–2025)

ABSTRACT. The Dirac operator with MIT bag boundary condition in a bounded convex domain is shown to be always self-adjoint in the H^1 -setting. This allows one to show that such operators appear as limit of Dirac operators with large positive mass outside the domain. Similar results were previously known for smooth domains only.

INTRODUCTION

For $n \geq 2$ and $N := 2^{\lfloor \frac{n+1}{2} \rfloor}$ let $\alpha_1, \dots, \alpha_n, \beta$ be pairwise anticommuting Hermitian $N \times N$ matrices with $\alpha_k^2 = \beta^2 = I_N$, where I_N is the $N \times N$ identity matrix (see Appendix A for a possible explicit choice). The Euclidean Dirac operator D_m with a mass $m \in \mathbb{R}$ acts on vector functions $f : \mathbb{R}^n \rightarrow \mathbb{C}^N$ by the differential expression

$$D_m f := -i \sum_{k=1}^n \alpha_k \partial_k f + m \beta f, \quad (1)$$

and it is a central object in the relativistic quantum mechanics, see e.g. [37].

Now let $\Omega \subset \mathbb{R}^n$ be a bounded domain with sufficiently regular (e.g. Lipschitz) boundary and outer unit normal ν . By the MIT bag operator with mass m in Ω one means the operator A_m^Ω acting in $L^2(\Omega, \mathbb{C}^N)$ as $f \mapsto D_m f$ on the functions f satisfying the MIT bag boundary condition $f = -i\beta\alpha \cdot \nu f$ on $\partial\Omega$, where we use the writing

$$\alpha \cdot x = \sum_{k=1}^n x_k \alpha_k \text{ for any } x = (x_1, \dots, x_n) \in \mathbb{R}^n$$

and denote a function on Ω and its trace on $\partial\Omega$ by the same symbol for the sake of readability. The precise regularity of the functions f in the operator domain guaranteeing “good” properties of A_m^Ω will be discussed below in greater detail.

The Dirac operator with the above boundary condition appeared initially in the physics literature [13, 17, 24], in particular, as a model of quark confinement in hadrons. One of the interesting features observed is that the MIT bag operator appears as the limit of Dirac operators in the whole space with a large mass applied outside Ω [12], which explains the fact that the MIT bag boundary condition is often referred to as the “infinite mass boundary condition” (and it is usually considered as an analog of the Dirichlet boundary condition for the Laplacian). While the study in the physics literature has a long story, it seems that the mathematically rigorous analysis of the operator A_m^Ω only became active in the last decade. For Ω with smooth boundary and $n \in \{2, 3\}$ several authors [3, 7, 9, 31] have shown that the above operator A_m^Ω is self-adjoint if considered on the domain

$$\{f \in H^1(\Omega, \mathbb{C}^N) : f = -i\beta\alpha \cdot \nu f \text{ on } \partial\Omega\}. \quad (2)$$

2020 *Mathematics Subject Classification.* Primary: 35P20, Secondary: 81Q05, 35F45, 47B25.

Key words and phrases. Dirac operator, boundary condition, convex domain, self-adjointness.

As noted in [30], the result holds in any dimension as a corollary of more general results on manifolds with boundaries [5, 6]. Moreover, if one denotes by $E_j(T)$ the j -th eigenvalue of a lower semibounded operator T (assuming the usual numbering in the non-decreasing order with multiplicities counted), for each $j \in \mathbb{N}$ one has

$$E_j((A_m^\Omega)^2) = \lim_{M \rightarrow +\infty} E_j((B_{m,M}^\Omega)^2), \quad (3)$$

where $B_{m,M}^\Omega$ is the Dirac operator in $L^2(\mathbb{R}^n, \mathbb{C}^N)$ with mass m in Ω and mass M in Ω^c ,

$$\begin{aligned} B_{m,M}^\Omega : f &\mapsto D_0 f + (m\mathbb{1}_\Omega + M\mathbb{1}_{\Omega^c})\beta f \equiv D_m f + (M - m)\mathbb{1}_{\Omega^c}\beta f, \\ \text{dom } B_{m,M}^\Omega &= H^1(\mathbb{R}^n, \mathbb{C}^N), \end{aligned} \quad (4)$$

see [36] for $n = 2$, or [2] for $n = 3$, or [30] for arbitrary n . The relation (3) is usually viewed as the mathematical expression of the infinite mass limit mentioned above. Note that there are some extensions to unbounded domains [4] and spin manifolds [20], and some estimates for the rate of convergence are available as well [2, 11]. It should be noted that the proofs of the convergence are mainly based on the expression

$$\|A_m^\Omega f\|_{L^2(\Omega, \mathbb{C}^N)}^2 = \int_\Omega (|\nabla f|^2 + m^2|f|^2) dx + \int_{\partial\Omega} \left(m + \frac{H}{2}\right) |f|^2 d\mathcal{H}^{n-1}, \quad f \in \text{dom } A_m^\Omega, \quad (5)$$

where $H : x \mapsto \text{tr d}|_x \nu$ is the mean curvature of $\partial\Omega$ and $\mathcal{H}^{n-1}$ is the $(n-1)$ -dimensional Hausdorff measure, and on the use of tubular coordinates near $\partial\Omega$. Remark that the same expression also plays a central role in the spectral optimization [1].

While there is a vast literature on boundary value problems for Dirac-type operators on domains/manifolds with smooth boundaries, see e.g. [5, 6, 14, 15, 22] and references therein, the case of non-smooth boundaries is much less elaborated. The paper [28] studied the self-adjointness of A_m^Ω for the case $n = 2$ when Ω is a polygon, and it was shown that the domain (2) leads to a self-adjoint operator if and only if the polygon is convex (more generally, the deficiency indices coincide with the number of concave corners). In [16] it was shown that A_m^Ω is self-adjoint on the domain (2) if Ω is a convex circular cone in three dimensions. As shown in [8, 10, 33], for bounded Lipschitz Ω and $n \in \{2, 3\}$ the operator A_m^Ω becomes self-adjoint if considered on the larger domain

$$\left\{ f \in H^{\frac{1}{2}}(\Omega, \mathbb{C}^N) : D_m f \in L^2(\Omega, \mathbb{C}^N), f = -i\beta\alpha \cdot \nu f \text{ on } \partial\Omega \right\},$$

and the paper [38] studied the local regularity of eigenfunctions. It should be noted that the lost of regularity also destroys the proofs for (3), as the finiteness of the summands in the crucially important identity (5) implicitly means the inclusion $\text{dom } A_m^\Omega \subset H^1(\Omega, \mathbb{C}^N)$. The paper [11] contains an alternative proof of (3) for $n = 3$ and smooth Ω with the help of the resolvents, but its main constructions are based on the microlocal analysis, and no obvious extension to non-smooth domains is expected. Overall, to our knowledge, there are no proofs of the asymptotics (3) for any non-smooth Ω .

The main goal of the present paper is to show that the above results for smooth Ω can be extended to a large class of non-smooth but *convex* Ω . Remark that the convexity is one of the standard assumptions guaranteeing maximal regularity for Laplacians with various boundary conditions, see e.g. [21, Chap. 3] for most classical results or [25, 26] for recent developments. Before passing to rigorous formulations we recall that any bounded convex domain has Lipschitz boundary [21, Corollary 1.2.2.3], hence, the usual machineries in Sobolev spaces (embeddings, traces, extension operators) are available. Our central result is as follows:

Theorem 1 (Self-adjointness). *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with outer unit normal ν (which is defined a.e. on $\partial\Omega$), and let $m \in \mathbb{R}$. Then the operator*

$$A_m^\Omega : f \mapsto D_m f, \quad \text{dom } A_m^\Omega = \left\{ f \in H^1(\Omega, \mathbb{C}^N) : f = -i\beta\alpha \cdot \nu f \text{ on } \partial\Omega \right\}, \quad (6)$$

is self-adjoint in $L^2(\Omega, \mathbb{C}^N)$ with compact resolvent.

For the subsequent discussion we impose a better regularity of the boundary.

Definition 2. Let $U \subset \mathbb{R}^n$ be a bounded domain with Lipschitz boundary. A point $s \in \partial U$ is called *regular* if U coincides with a C^∞ -smooth domain in an open neighborhood of s , otherwise it is called *singular*. One denotes

$$\begin{aligned} \partial_\infty U &:= \{s \in \partial U : s \text{ is regular}\} && (\text{the regular boundary of } U), \\ \partial_0 U &:= \{s \in \partial U : s \text{ is singular}\} && (\text{the singular boundary of } U). \end{aligned}$$

The domain U will be called *piecewise smooth* if its singular boundary is contained in the union of finitely many compact $(n-2)$ -dimensional submanifolds of $\mathbb{R}^n$.

We show:

Theorem 3 (Quadratic form). *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary and outer unit normal ν (which is defined at least on the regular boundary), then for any $f \in \text{dom } A_m^\Omega$ one has*

$$\|A_m^\Omega f\|_{L^2(\Omega, \mathbb{C}^N)}^2 = \int_\Omega \left(|\nabla f|^2 + m^2 |f|^2 \right) dx + \int_{\partial\Omega} \left(m + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1},$$

where $H : x \mapsto \text{tr } d|_x \nu \geq 0$ is the mean curvature defined on the regular boundary of Ω .

By imposing an additional geometric condition we are finally able to extend (3) to a large class of non-smooth domains:

Theorem 4 (Infinite mass limit). *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary, and assume that the mean curvature of the regular boundary is bounded. Then for any $m \in \mathbb{R}$ and $j \in \mathbb{N}$ one has $E_j((A_m^\Omega)^2) = \lim_{M \rightarrow +\infty} E_j((B_{m,M}^\Omega)^2)$, where the operator $B_{m,M}^\Omega$ acting in $L^2(\mathbb{R}^n, \mathbb{C}^N)$ is defined by (4).*

The assumptions of Theorem 4 are satisfied, in particular, if Ω is a bounded convex polyhedron (i.e. a bounded domain obtained as an intersection of halfspaces), or a convex curvilinear polyhedron (i.e. obtained as a diffeomorphic image of a bounded convex polyhedron), or is a convex piecewise smooth domain obtained as the intersection of finitely many C^∞ -smooth bounded domains. On the other hand, the assumption on the boundedness of the mean curvature excludes e.g. the three-dimensional domains whose boundary has conical singularities of the form $x_3 = \sqrt{x_1^2 + x_2^2}$. We believe that the assumption on the mean curvature is purely technical, see Remark 20 at the end of the paper.

Our proof of Theorem 1 is presented in Section 1, and it is based on adaptations of some constructions in Grisvard's book [21, Sec. 3.2] for second-order operators. Remark 10 indicates possible extensions. In Section 2 we derive some approximation results for Sobolev spaces. This allows one to show the essential self-adjointness of A_m^Ω on smooth functions vanishing near the singular boundary (Lemma 16), which is an essential ingredient for the subsequent analysis. In particular, it is used in Section 3 to prove Theorem 3 by transferring the results known for the smooth case. Theorem 4 is shown in Section 4, and is mainly in the spirit of the analysis of [30], while a more involved choice of test functions is used, and the constructions with tubular

coordinates near $\partial\Omega$ (which are now unavailable due to non-smoothness) are replaced by some estimates for Robin Laplacians in exterior domains. The short Appendix A collects a necessary information on the Dirac matrices.

1. PROOF OF THEOREM 1 (SELF-ADJOINTNESS)

As a starting point let us summarize rigorously some results on smooth domains, see [30, Lem. 2.1 and Prop. A.2].

Lemma 5. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with C^∞ -smooth boundary and outer unit normal ν , and let $m \in \mathbb{R}$. Then the operator A_m^Ω defined by (6) is self-adjoint in $L^2(\Omega, \mathbb{C}^N)$, and for any $f \in \text{dom } A_m^\Omega$ one has the identity (5).*

As the parameter m only results in adding a bounded symmetric perturbation, it is sufficient to show that the operator

$$A := A_0^\Omega$$

is self-adjoint and has compact resolvent. The integration by parts show that A is symmetric, and its self-adjointness is equivalent to the equalities $\text{ran}(A \pm i) = L^2(\Omega, \mathbb{C}^N)$. We will show $\text{ran}(A + i) = L^2(\Omega, \mathbb{C}^N)$, as the other condition is proved completely analogously by adjusting the signs.

Our constructions will be based on the following results of convex analysis, see e.g. [34, Sec. 3.4] and [21, Lemma 3.2.3.2]:

Lemma 6. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain. There exist bounded convex C^∞ -smooth domains $\Omega_p \subset \mathbb{R}^n$, $p \in \mathbb{N}$, such that $\Omega \subset \Omega_p$ for all p and $d_H(\partial\Omega_p, \partial\Omega) \xrightarrow{p \rightarrow \infty} 0$, where d_H stands for the Hausdorff distance. In addition, one can find open subsets $V_k \subset \mathbb{R}^n$, $k \in \{1, \dots, K\}$, with the following properties:*

- for each k there exist Cartesian coordinates $y_1^k, \dots, y_n^k$ in which V_k is a hypercube,

$$V_k = \left\{ (y_1^k, \dots, y_n^k) : -a_j^k < y_j^k < a_j^k \text{ for all } j \in \{1, \dots, n\} \right\}, \quad a_j^k > 0,$$

- for all $p \in \mathbb{N}$ and each $k \in \{1, \dots, K\}$ there exist a Lipschitz function h^k and a C^∞ -smooth function h_p^k defined on

$$V'_k = \left\{ (y_1^k, \dots, y_{n-1}^k) : -a_j^k < y_j^k < a_j^k \text{ for all } j \in \{1, \dots, n-1\} \right\}$$

such that

$$h^k(z^k) \leq h_p^k(z^k), \quad |h^k(z^k)| \leq \frac{a_n^k}{2}, \quad |h_p^k(z^k)| \leq \frac{a_n^k}{2} \text{ for all } z^k \in V'_k,$$

$$\Omega \cap V_k = \left\{ y^k = (z^k, y_n^k) \in V_k : z^k \in V'_k, y_n^k < h^k(z^k) \right\},$$

$$\Omega_p \cap V_k = \left\{ y^k = (z^k, y_n^k) \in V_k : z^k \in V'_k, y_n^k < h_p^k(z^k) \right\},$$

$$\partial\Omega \cap V_k = \left\{ y^k = (z^k, y_n^k) \in V_k : z^k \in V'_k, y_n^k = h^k(z^k) \right\},$$

$$\partial\Omega_p \cap V_k = \left\{ y^k = (z^k, y_n^k) \in V_k : z^k \in V'_k, y_n^k = h_p^k(z^k) \right\},$$

- $\partial\Omega \subset \bigcup_{k=1}^K V_k$ and $\partial\Omega_p \subset \bigcup_{k=1}^K V_k$ for all $p \in \mathbb{N}$,
- for each $k \in \{1, \dots, K\}$ one has $h_p^k \xrightarrow{p \rightarrow \infty} h^k$ uniformly in V'_k ,
- there is an $L > 0$ such that $|\nabla h^k(z^k)| \leq L$ and $|\nabla h_p^k(z^k)| \leq L$ for all a.e. $z^k \in V'_k$, all $k \in \{1, \dots, K\}$ and all $p \in \mathbb{N}$,
- for each $k \in \{1, \dots, K\}$ one has $\nabla h_p^k(z^k) \xrightarrow{p \rightarrow \infty} \nabla h^k(z^k)$ for a.e. $z^k \in V'_k$.

Let us approximate Ω by bounded convex C^∞ -smooth domains Ω_p (with $p \in \mathbb{N}$) as in Lemma 6 and consider the associated Dirac operators

$$\tilde{A}_p := A_0^{\Omega_p},$$

which are self-adjoint in $L^2(\Omega_p, \mathbb{C}^N)$ by Lemma 5.

Let $g \in L^2(\Omega, \mathbb{C}^N)$. For each p denote by $\tilde{g}_p$ the continuation of g by zero to Ω_p then due to the self-adjointness of $\tilde{A}_p$ there is a unique $\tilde{f}_p \in \text{dom } \tilde{A}_p$ with $(\tilde{A}_p + i)\tilde{f}_p = \tilde{g}_p$. Let H_p be the mean curvature of $\partial\Omega_p$, then $H_p \geq 0$ due to convexity of Ω_p , and by Lemma 5 we obtain

$$\begin{aligned} \|g\|_{L^2(\Omega, \mathbb{C}^N)}^2 &= \|\tilde{g}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 = \|(\tilde{A}_p + i)\tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 = \|\tilde{A}_p \tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 + \|\tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 \\ &= \int_{\Omega_p} (|\nabla \tilde{f}_p|^2 + |\tilde{f}_p|^2) dx + \frac{1}{2} \int_{\partial\Omega_p} H_p |\tilde{f}_p|^2 d\mathcal{H}^{n-1} \\ &\geq \int_{\Omega_p} (|\nabla \tilde{f}_p|^2 + |\tilde{f}_p|^2) dx = \|\tilde{f}_p\|_{H^1(\Omega_p, \mathbb{C}^N)}^2. \end{aligned} \quad (7)$$

Denote by f_p the restriction of $\tilde{f}_p$ on Ω , then

$$\|f_p\|_{H^1(\Omega, \mathbb{C}^N)} \leq \|\tilde{f}_p\|_{H^1(\Omega_p, \mathbb{C}^N)} \leq \|g\|_{L^2(\Omega, \mathbb{C}^N)}, \quad (8)$$

which shows that the sequence $(f_p)_{p \in \mathbb{N}}$ is bounded in $H^1(\Omega, \mathbb{C}^N)$ and, hence, contains a weakly convergent subsequence. To keep simple notation we assume (by replacing the initial sequence of Ω_p by the corresponding subsequence) that $f_p \xrightarrow{p \rightarrow \infty} f$ weakly in $H^1(\Omega, \mathbb{C}^N)$ for some $f \in H^1(\Omega, \mathbb{C}^N)$. We are going to show that $f \in \text{dom } A$ with $(A + i)f = g$, which will conclude the proof of self-adjointness for A .

Due to the compactness of the embedding $H^1(\Omega, \mathbb{C}^N) \hookrightarrow L^2(\Omega, \mathbb{C}^N)$ we have

$$\|f_p - f\|_{L^2(\Omega, \mathbb{C}^N)} \xrightarrow{p \rightarrow \infty} 0. \quad (9)$$

Let $\varphi \in C_c^\infty(\Omega, \mathbb{C}^N)$ and $\tilde{\varphi}_p$ be its extension by zero on Ω_p . Then $\tilde{\varphi}_p \in C_c^\infty(\Omega_p, \mathbb{C}^N) \subset \text{dom } \tilde{A}_p$, and

$$\begin{aligned} \langle g, \varphi \rangle_{L^2(\Omega, \mathbb{C}^N)} &= \langle \tilde{g}_p, \tilde{\varphi}_p \rangle_{L^2(\Omega_p, \mathbb{C}^N)} = \langle (\tilde{A}_p + i)\tilde{f}_p, \tilde{\varphi}_p \rangle_{L^2(\Omega_p, \mathbb{C}^N)} \\ &= \langle \tilde{f}_p, (\tilde{A}_p - i)\tilde{\varphi}_p \rangle_{L^2(\Omega_p, \mathbb{C}^N)} = \langle \tilde{f}_p, (D_0 - i)\tilde{\varphi}_p \rangle_{L^2(\Omega_p, \mathbb{C}^N)} \\ &= \langle f_p, (D_0 - i)\varphi \rangle_{L^2(\Omega, \mathbb{C}^N)} \xrightarrow[p \rightarrow \infty]{\text{by (9)}} \langle f, (D_0 - i)\varphi \rangle_{L^2(\Omega, \mathbb{C}^N)}. \end{aligned}$$

Hence,

$$\int_{\Omega} \langle g(x), \varphi(x) \rangle_{\mathbb{C}^N} dx = \int_{\Omega} \langle f(x), (D_0 - i)\varphi(x) \rangle_{\mathbb{C}^N} dx \text{ for all } \varphi \in C_c^\infty(\Omega, \mathbb{C}^N),$$

which means $(D_0 + i)f = g$ in $\mathcal{D}'(\Omega, \mathbb{C}^N)$. As $f \in H^1(\Omega, \mathbb{C}^N)$ by construction, it remains to check that f satisfies the required boundary condition.

Let V_k, V'_k, h^k, h_p^k and L be as in Lemma 6, and let $\nu^k(z^k)$ be the outer unit normal for Ω at the point $(z^k, h^k(z^k)) \in V_k \cap \partial\Omega$ and $\nu_p^k(z^k)$ be the outer unit normal for Ω_p at the point $(z^k, h_p^k(z^k)) \in V_k \cap \partial\Omega_p$, i.e.

$$\nu^k(z^k) = \frac{(-\nabla h^k(z^k), 1)}{\sqrt{1 + |\nabla h^k(z^k)|^2}}, \quad \nu_p^k(z^k) = \frac{(-\nabla h_p^k(z^k), 1)}{\sqrt{1 + |\nabla h_p^k(z^k)|^2}}.$$

In order to show that the function f satisfies the boundary condition we need to check that for any $k \in \{1, \dots, K\}$ one has

$$\int_{V_k \cap \partial\Omega} |(1 + i\beta\alpha \cdot \nu)f|^2 d\mathcal{H}^{n-1} \equiv \int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu^k(z^k))f(z^k, h^k(z^k)) \right|^2 \sqrt{1 + |\nabla h^k(z^k)|^2} dz^k = 0,$$

and in view of the bound $|\nabla h^k(z^k)| \leq L$ this is equivalent to

$$\int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu^k(z^k))f(z^k, h^k(z^k)) \right|^2 dz^k = 0. \quad (10)$$

By construction, the functions $\tilde{f}_p$ satisfy $(1 + i\beta\alpha \cdot \nu_p)\tilde{f}_p = 0$ on $\partial\Omega_p$, with ν_p being the unit normal for Ω_p , which means that for any $k \in \{1, \dots, K\}$ it holds that

$$\int_{V_k \cap \partial\Omega_p} |(1 + i\beta\alpha \cdot \nu_p)\tilde{f}_p|^2 d\mathcal{H}^{n-1} = 0,$$

hence,

$$\int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu_p^k(z^k))\tilde{f}_p(z^k, h^k(z^k)) \right|^2 dz^k = 0. \quad (11)$$

Our goal is to pass to the limit $p \rightarrow \infty$ in this last formula in order to obtain the required identities (10). As a preparation let us derive some preliminary estimates.

Lemma 7. *For any $F \in H^1(\mathbb{R}^n, \mathbb{C}^N)$ any $k \in \{1, \dots, K\}$ one has*

$$\int_{V'_k} \left| F(z^k, h_p^k(z^k)) - F(z^k, h^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0.$$

Proof. Using the Cauchy-Schwarz inequality we have

$$\begin{aligned} \int_{V'_k} \left| F(z^k, h_p^k(z^k)) - F(z^k, h^k(z^k)) \right|^2 dz^k &= \int_{V'_k} \left| \int_{h^k(z^k)}^{h_p^k(z^k)} \partial_{y_n^k} F(z^k, t) dt \right|^2 dz^k \\ &\leq \int_{V'_k} |h_p^k(z^k) - h^k(z^k)| \int_{h^k(z^k)}^{h_p^k(z^k)} |\partial_{y_n^k} F(z^k, t)|^2 dt dz^k \\ &\leq \sup_{z^k \in V'_k} |h_p^k(z^k) - h^k(z^k)| \int_{V_k \cap \Omega_p} |\partial_{y_n^k} F|^2 dy^k \\ &\leq \sup_{z^k \in V'_k} |h_p^k(z^k) - h^k(z^k)| \|F\|_{H^1(\mathbb{R}^n, \mathbb{C}^N)}^2 \xrightarrow{p \rightarrow \infty} 0 \end{aligned}$$

due to the uniform convergence of h_p^k to h^k on V'_k . $\square$

Lemma 8. *Let $F \in H^1(\mathbb{R}^n, \mathbb{C}^N)$ be a continuation of f on $\mathbb{R}^n$, then for any $k \in \{1, \dots, K\}$ one has*

$$\int_{V'_k} \left| F(z^k, h_p^k(z^k)) - \tilde{f}_p(z^k, h_p^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0.$$

Proof. We need to show that

$$\int_{V'_k} \left| \eta_p(z^k, h_p^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0 \text{ for the functions } \eta_p := F - \tilde{f}_p \in H^1(\Omega_p).$$

First remark that due to (8) one can find a $c > 0$ such that

$$\|\eta_p\|_{H^1(\Omega_p, \mathbb{C}^N)} \leq c \text{ for all } p \in \mathbb{N}. \quad (12)$$

Furthermore, by assumption $\eta_p \rightarrow 0$ weakly in $H^1(\Omega)$. Using the boundedness of the trace operator $H^1(\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$ and the compactness of the embedding $H^{\frac{1}{2}}(\partial\Omega) \hookrightarrow L^2(\partial\Omega)$ we obtain

$$\int_{\partial\Omega} |\eta_p|^2 d\mathcal{H}^{n-1} \xrightarrow{p \rightarrow \infty} 0,$$

and by combining with

$$\begin{aligned} \int_{V'_k} \left| \eta_p(z^k, h^k(z^k)) \right|^2 dz^k &\leq \frac{1}{\sqrt{1+L^2}} \int_{V'_k} \left| \eta_p(z^k, h^k(z^k)) \right|^2 \sqrt{1+|\nabla h^k(z^k)|^2} dz^k \\ &= \frac{1}{\sqrt{1+L^2}} \int_{V_k \cap \partial\Omega} |\eta_p|^2 d\mathcal{H}^{n-1} \leq \frac{1}{\sqrt{1+L^2}} \int_{\partial\Omega} |\eta_p|^2 d\mathcal{H}^{n-1} \end{aligned}$$

we conclude that

$$\int_{V'_k} \left| \eta_p(z^k, h^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0. \quad (13)$$

Now we estimate

$$\begin{aligned} \int_{V'_k} \left| \eta_p(z^k, h_p^k(z^k)) \right|^2 dz^k &= \int_{V'_k} \left| \eta_p(z^k, h^k(z^k)) + \int_{h^k(z^k)}^{h_p^k(z^k)} \partial_{y_n^k} \eta_p(z^k, t) dt \right|^2 dz^k \\ &\leq 2 \int_{V'_k} \left| \eta_p(z^k, h^k(z^k)) \right|^2 dz^k + 2 \int_{V'_k} \left| \int_{h^k(z^k)}^{h_p^k(z^k)} \partial_{y_n^k} \eta_p(z^k, t) dt \right|^2 dz^k. \end{aligned}$$

For $p \rightarrow \infty$ the first summand on the right-hand side converges to zero by (13), and the second summand is easily controlled using Cauchy-Schwarz inequality:

$$\begin{aligned} \int_{V'_k} \left| \int_{h^k(z^k)}^{h_p^k(z^k)} \partial_{y_n^k} \eta_p(z^k, t) dt \right|^2 dz^k &\leq \int_{V'_k} |h_p^k(z^k) - h^k(z^k)| \int_{h^k(z^k)}^{h_p^k(z^k)} |\partial_{y_n^k} \eta_p(z^k, t)|^2 dt dz^k \\ &\leq \sup_{z^k \in V'_k} |h_p^k(z^k) - h^k(z^k)| \int_{V_k \cap \Omega_p} |\partial_{y_n^k} \eta_p|^2 dy^k \\ &\leq \sup_{z^k \in V'_k} |h_p^k(z^k) - h^k(z^k)| \|\eta_p\|_{H^1(V_k \cap \Omega_p)}^2 \\ \text{use (12): } &\leq c^2 \sup_{z^k \in V'_k} |h_p^k(z^k) - h^k(z^k)| \xrightarrow{p \rightarrow \infty} 0 \end{aligned}$$

due to the uniform convergence of h_p^k to h^k on V'_k . □

Let $k \in \{1, \dots, K\}$ and let $F \in H^1(\mathbb{R}^n, \mathbb{C}^N)$ be an arbitrary continuation of f , then

$$\|\beta\alpha \cdot \nu^k(z^k)\| \leq 1 \text{ and } \|\beta\alpha \cdot \nu_p^k(z^k)\| \leq 1 \text{ for all } p \in \mathbb{N} \text{ and a.e. } z^k \in V'_k, \quad (14)$$

$$\beta\alpha \cdot \nu_p^k(z^k) \xrightarrow{p \rightarrow \infty} \beta\alpha \cdot \nu^k(z^k) \text{ for a.e. } z^k \in V'_k. \quad (15)$$

Using the representation

$$\begin{aligned} (1 + i\beta\alpha \cdot \nu^k(z^k)) f(z^k, h^k(z^k)) &= (1 + i\beta\alpha \cdot \nu^k(z^k)) F(z^k, h^k(z^k)) \\ &= (i\beta\alpha \cdot \nu^k(z^k) - i\beta\alpha \cdot \nu_p^k(z^k)) F(z^k, h^k(z^k)) \\ &\quad + (1 + i\beta\alpha \cdot \nu_p^k(z^k)) (F(z^k, h^k(z^k)) - F(z^k, h_p^k(z^k))) \\ &\quad + (1 + i\beta\alpha \cdot \nu_p^k(z^k)) (F(z^k, h_p^k(z^k)) - \tilde{f}_p(z^k, h_p^k(z^k))) \\ &\quad + (1 + i\beta\alpha \cdot \nu_p^k(z^k)) \tilde{f}_p(z^k, h_p^k(z^k)) \end{aligned}$$

we obtain

$$\begin{aligned} \int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu^k(z^k)) f(z^k, h^k(z^k)) \right|^2 dz^k &\leq 4(I_p^1 + I_p^2 + I_p^3 + I_p^4) \\ \text{for } I_p^1 &:= \int_{V'_k} \left| (i\beta\alpha \cdot \nu^k(z^k) - i\beta\alpha \cdot \nu_p^k(z^k)) F(z^k, h^k(z^k)) \right|^2 dz^k, \end{aligned} \quad (16)$$

$$\begin{aligned}
I_p^2 &:= \int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu_p^k(z^k)) \left(F(z^k, h^k(z^k)) - F(z^k, h_p^k(z^k)) \right) \right|^2 dz^k, \\
I_p^3 &:= \int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu_p^k(z^k)) \left(F(z^k, h_p^k(z^k)) - \tilde{f}_p(z^k, h_p^k(z^k)) \right) \right|^2 dz^k, \\
I_p^4 &:= \int_{V'_k} \left| (1 + i\beta\alpha \cdot \nu_p^k(z^k)) \tilde{f}_p(z^k, h_p^k(z^k)) \right|^2 dz^k \stackrel{(11)}{=} 0.
\end{aligned}$$

In virtue of (14) and (15), the dominated convergence theorem implies $I_p^1 \xrightarrow{p \rightarrow \infty} 0$. Using (14) again we estimate

$$\begin{aligned}
I_p^2 &\leq 4 \int_{V'_k} \left| F(z^k, h^k(z^k)) - F(z^k, h_p^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0 \text{ by Lemma 7,} \\
I_p^3 &\leq 4 \int_{V'_k} \left| F(z^k, h_p^k(z^k)) - \tilde{f}_p(z^k, h_p^k(z^k)) \right|^2 dz^k \xrightarrow{p \rightarrow \infty} 0 \text{ by Lemma 8.}
\end{aligned}$$

This shows that the right-hand side of (16) converges to 0 for $p \rightarrow \infty$, hence, the left-hand side is zero. The proof of self-adjointness is completed.

For the sake of completeness we include the following standard assertion:

Lemma 9. *The graph norm of A on $\text{dom } A$ is equivalent to the H^1 -norm, and A has compact resolvent.*

Proof. The self-adjointness of A implies that $\text{dom } A$ is complete with respect to the graph norm. At the same time, $\text{dom } A$ is also complete with respect to the H^1 -norm being the kernel of the bounded linear operator $H^1(\mathbb{R}^n, \mathbb{C}^N) \ni f \mapsto (1 + i\beta\alpha \cdot \nu)f|_{\partial\Omega} \in L^2(\partial\Omega)$. The embedding operator $(\text{dom } A \text{ with } H^1\text{-norm}) \ni f \mapsto f \in (\text{dom } A \text{ with the graph norm})$ is obviously bijective and bounded, hence, its inverse is bounded due to the closed graph theorem, and this shows that the both norms are equivalent. It follows that $\text{dom } A$ with the graph norm is continuously embedded into $H^1(\Omega, \mathbb{C}^N)$ and, therefore, compactly embedded in $L^2(\Omega, \mathbb{C}^N)$, which gives the conclusion. $\square$

Remark 10. One easily sees that the convexity was actually used only to establish a uniform upper bound for the H^1 -norms of the functions $\tilde{f}_p$ in (7). In fact, the proof can be adapted to a larger class of domains of Ω as follows. Let us drop the convexity assumption but require instead that Ω can be approximated (in the sense described in Lemma 6) by C^∞ -smooth domains Ω_p such that the mean curvatures H_p of their boundaries are uniformly bounded from below, $H_p \geq -2m$ for some $m > 0$. Then one can consider the operators $A := A_m^\Omega$ and $\tilde{A}_p := A_m^{\Omega_p}$. If $g \in L^2(\Omega, \mathbb{C}^N)$ and $\tilde{g}_p$ are its extensions by zero to Ω_p , then we can find $\tilde{f}_p \in \text{dom } \tilde{A}_p$ with $(\tilde{A}_p + i)\tilde{f}_p = \tilde{g}_p$, and (7) can be replaced by

$$\begin{aligned}
\|g\|_{L^2(\Omega, \mathbb{C}^N)}^2 &= \|\tilde{g}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 = \|(\tilde{A}_p + i)\tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 = \|\tilde{A}_p \tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 + \|\tilde{f}_p\|_{L^2(\Omega_p, \mathbb{C}^N)}^2 \\
&= \int_{\Omega_p} (|\nabla \tilde{f}_p|^2 + (m^2 + 1)|\tilde{f}_p|^2) dx + \int_{\partial\Omega_p} \left(m + \frac{H_p}{2} \right) |\tilde{f}_p|^2 d\mathcal{H}^{n-1} \\
&\geq \int_{\Omega_p} (|\nabla \tilde{f}_p|^2 + |\tilde{f}_p|^2) dx = \|\tilde{f}_p\|_{H^1(\Omega_p, \mathbb{C}^N)}^2,
\end{aligned}$$

and the rest of the proof remains literally the same, which provides the self-adjointness of A for such Ω . Simple geometric considerations show that for $n = 2$ these observations apply, for example, to curvilinear polygons without concave corners.

2. APPROXIMATIONS

In this section we establish several approximation results for Sobolev spaces on piecewise smooth domains. The constructions are inspired by [18]. We need first a simple result on the approximation of non-smooth domains by smooth ones.

Lemma 11 (“Rounding the corners”). *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain, then for any open neighborhood $V_0 \subset \mathbb{R}^n$ of its singular boundary $\partial_0\Omega$ one can find a bounded domain $\Omega_0 \subset \mathbb{R}^n$ with C^∞ -smooth boundary such that Ω and Ω_0 coincide outside V_0 .*

Proof. Without loss of generality assume that $0 \in \Omega$, then due to convexity we can describe Ω using polar coordinates,

$$\Omega = \{r\theta : \theta \in \mathbb{S}^{n-1}, 0 \leq r < R(\theta)\}, \quad \partial\Omega = \{R(\theta)\theta : \theta \in \mathbb{S}^{n-1}\},$$

with a Lipschitz function $R : \mathbb{S}^{n-1} \rightarrow (0, \infty)$ which is C^∞ -smooth on the open set

$$S_\infty := \{\theta \in \mathbb{S}^{n-1} : R(\theta)\theta \in \partial_\infty\Omega\} \subset \mathbb{S}^{n-1}.$$

Remark $\partial_0\Omega$ is compact and that the Lipschitz map

$$\Pi : \mathbb{R}^n \setminus \{0\} \ni x \mapsto \frac{x}{|x|} \in \mathbb{S}^{n-1}$$

is the inverse of $\theta \mapsto R(\theta)\theta$, which means that $S_0 := \mathbb{S}^{n-1} \setminus S_\infty \equiv \Pi(\partial_0\Omega)$ is a compact subset of the open set $\Theta_0 := \Pi(V_0) \subset \mathbb{S}^{n-1}$, and $\{R(\theta)\theta : \theta \in S_0\} = \partial_0\Omega \subset V_0$.

We now choose two open neighborhoods Θ_1, Θ_2 of S_0 in $\mathbb{S}^{n-1}$ such that $\overline{\Theta_2} \subset \Theta_1$ and $\overline{\Theta_1} \subset \Theta_0$. Due to compactness of $\overline{\Theta_1}$ there is an $\varepsilon > 0$ such that

$$\{r\theta : \theta \in \overline{\Theta_1}, |r - R(\theta)| \leq \varepsilon\} \subset V_0,$$

and we can choose a C^∞ -function $R_1 : \Theta_1 \rightarrow (0, \infty)$ such that $|R_1(\theta) - R(\theta)| \leq \varepsilon$ for all $\theta \in \Theta_1$. Now choose a cut-off function $\chi \in C^\infty(\mathbb{S}^{n-1})$ with $0 \leq \chi \leq 1$ such that $\text{supp } \chi \subset \Theta_1$ and $\chi = 1$ on Θ_2 , then $R_0 := (1 - \chi)R + \chi R_1 : \mathbb{S}^{n-1} \rightarrow (0, \infty)$ is smooth, and the graphs of $\theta \mapsto R(\theta)\theta$ and $\theta \mapsto R_0(\theta)\theta$ coincide outside V_0 . It follows that the domain

$$\Omega_0 := \{r\theta : \theta \in \mathbb{S}^{n-1}, 0 \leq r < R_0(\theta)\}$$

satisfies all the requirements. $\square$

The following assertion is [18, Lem. 2.4] with a reformulation adapted to the subsequent use:

Lemma 12. *Let $\chi \in C_c^\infty(\mathbb{R})$ with $0 \leq \chi \leq 1$ such that $\text{supp } \chi \subset (-1, 1)$ and $\chi = 1$ in an open neighborhood of 0. For $\alpha \in (0, 1)$ and $n \in \mathbb{N}$ define functions*

$$\rho_{\alpha,n} : [0, \infty) \ni t \mapsto \min\{t^\alpha, 1\}(1 - \chi(nt)) \in \mathbb{R}.$$

Then one can choose $\alpha_j \xrightarrow{j \rightarrow \infty} 0$ and $n_j \xrightarrow{j \rightarrow \infty} \infty$ such that the functions

$$\psi_j : \mathbb{R}^2 \ni x \mapsto \rho_{\alpha_j, n_j}(|x|) \in \mathbb{R}$$

satisfy:

- (a) *for any j one has $0 \leq \psi_j \leq 1$, the function ψ_j vanishes in an open neighborhood of 0 in $\mathbb{R}^2$, and $\psi_j(x) \xrightarrow{j \rightarrow \infty} 1$ for all $x \neq 0$,*
- (b) *$\|\psi_j f - f\|_{H^1(\mathbb{R}^2)} \xrightarrow{j \rightarrow \infty} 0$ for any $f \in H^1(\mathbb{R}^2)$,*
- (c) *there is a constant $C > 0$ such that $\|\psi_j f\|_{H^1(\mathbb{R}^2)} \leq C\|f\|_{H^1(\mathbb{R}^2)}$ for any $f \in H^1(\mathbb{R}^2)$ and any $j \in \mathbb{N}$.*

Remark that [18] only states (b) explicitly. The part (a) follows from the explicit structure of ψ_j , and (c) follows from (b) in virtue of Banach-Steinhaus theorem.

Lemma 13 (Cut-off near a submanifold). *Let $n \geq 2$ and Γ be a compact $(n-2)$ -dimensional submanifold in $\mathbb{R}^n$. For $x \in \mathbb{R}^n$ let $d(x)$ be the distance from x to Γ . Then the functions $\varphi_j := \psi_j \circ d$ with ψ_j from Lemma 12 satisfy*

$$\|\varphi_j f - f\|_{H^1(\mathbb{R}^n)} \xrightarrow{j \rightarrow \infty} 0 \text{ for any } f \in H^1(\mathbb{R}^n). \quad (17)$$

In addition, $\varphi_j = 0$ in an open neighborhood of Γ , with $0 \leq \varphi_j \leq 1$, and $\varphi_j(x) \xrightarrow{j \rightarrow \infty} 1$ for any $x \in \mathbb{R}^n \setminus \Gamma$.

Proof. For $r > 0$ denote $B_2(0, r) := \{x \in \mathbb{R}^2 : |x| < r\}$. It is a classical result of differential geometry that for some sufficiently small $\varepsilon > 0$ the ε -neighborhood Γ_ε of Γ admits a tubular neighborhood, see e.g. [27, Thm. 10.19]. In the coordinate form this means that one can cover Γ_ε by finitely many open sets $V_1, \dots, V_L$ such that for any $s \in \Gamma_\ell := V_\ell \cap \Gamma$ one can choose an orthonormal basis $(n_1(s), n_2(s))$ in the normal space to Γ at s depending smoothly on s and the maps

$$\Phi_\ell : \Gamma_\ell \times B_2(0, \varepsilon) \ni (s, t_1, t_2) \mapsto s + t_1 n_1(s) + t_2 n_2(s) \in V_\ell,$$

are diffeomorphisms, with $d(\Phi_\ell(s, t_1, t_2)) = \sqrt{t_1^2 + t_2^2} = |t|$, and the map $f \mapsto f \circ \Phi_\ell$ defines an isomorphism between $H^1(V_\ell)$ and $H^1(\Gamma_\ell \times B_2(0, \varepsilon))$. Without loss of generality we assume that $\varepsilon \in (0, 1)$.

Let $f \in H^1(\mathbb{R}^n)$. Note first that by construction one has $\varphi_j f = f$ outside Γ_1 . Furthermore, for $x \in \Gamma_1 \setminus \Gamma_\varepsilon$ and all sufficiently large j one has $(\varphi_j f)(x) = d(x)^{\alpha_j} f(x)$, and using $\alpha_j \rightarrow 0$ and the dominated convergence theorem we arrive at $\|\varphi_j f - f\|_{L^2(\Gamma_1 \setminus \Gamma_\varepsilon)} \xrightarrow{j \rightarrow \infty} 0$. In addition, for the same x one has

$$\nabla(\varphi_j f)(x) - \nabla f(x) = d(x)^{\alpha_j} \nabla f(x) - \nabla f(x) + \alpha_j \frac{\nabla d(x)}{d(x)^{1-\alpha_j}} f(x),$$

$$\int_{\Gamma_1 \setminus \Gamma_\varepsilon} |\nabla(\varphi_j f)(x) - \nabla f(x)|^2 dx \leq 2 \int_{\Gamma_1 \setminus \Gamma_\varepsilon} |d^{\alpha_j} \nabla f - \nabla f|^2 dx + 2\alpha_j^2 \int_{\Gamma_1 \setminus \Gamma_\varepsilon} \left| \frac{\nabla d}{d^{1-\alpha_j}} \right|^2 |f|^2 dx.$$

The first summand on the right-hand side of the last inequality converges to zero by the dominated convergence theorem, while for the second summand we have

$$2\alpha_j^2 \int_{\Gamma_1 \setminus \Gamma_\varepsilon} \left| \frac{\nabla d}{d^{1-\alpha_j}} \right|^2 |f|^2 dx \leq \frac{2\alpha_j^2}{\varepsilon^{2(1-\alpha_j)}} \int_{\Gamma_1 \setminus \Gamma_\varepsilon} |f|^2 dx \xrightarrow{j \rightarrow \infty} 0.$$

The above considerations show that $\|\varphi_j f - f\|_{H^1(\mathbb{R}^n \setminus \Gamma_\varepsilon)} \xrightarrow{j \rightarrow \infty} 0$, and it remains to check $\|\varphi_j f - f\|_{H^1(\Gamma_\varepsilon)} \xrightarrow{j \rightarrow \infty} 0$. In view of the preceding discussion it is sufficient to show that for each $\ell \in \{1, \dots, L\}$ one has

$$\hat{f}_j \xrightarrow{j \rightarrow \infty} \hat{f} \text{ in } H^1(\Gamma_\ell \times B_2(0, \varepsilon)), \quad \hat{f}_j := (\varphi_j f) \circ \Phi_\ell, \quad \hat{f} := f \circ \Phi_\ell.$$

One has $\hat{f}_j(s, t) = \psi_j(t) \hat{f}(s, t)$, in particular, $\nabla_s \hat{f}_j = \nabla_s \hat{f}$, and

$$\begin{aligned} \|\hat{f}_j - \hat{f}\|_{H^1(\Gamma_\ell \times B_2(0, \varepsilon_0))}^2 &= \int_{\Gamma_\ell} \left(\left\| \psi_j f(s, \cdot) - f(s, \cdot) \right\|_{H^1(B_2(0, \varepsilon))}^2 \right. \\ &\quad \left. + \left\| \psi_j \nabla_s f(s, \cdot) - \nabla_s f(s, \cdot) \right\|_{L^2(B_2(0, \varepsilon))}^2 \right) d\mathcal{H}^{n-2}(s). \end{aligned}$$

The subintegral expression admits integrable upper bound and converges to 0 for a.e. $s \in \Gamma_\ell$ as $j \rightarrow \infty$ by Lemma 12, so the whole integral converges to 0 for $j \rightarrow \infty$ by the dominated

convergence theorem. This completes the proof of (17), and the remaining claims on φ_j follow from the properties of ψ_j in Lemma 12. $\square$

Corollary 14. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with piecewise smooth boundary. Then one can find functions $\rho_j \in C^\infty(\mathbb{R}^n)$ with $j \in \mathbb{N}$ such that*

$$0 \leq \rho_j \leq 1, \quad \rho_j = 0 \text{ in an open neighborhood of } \partial_0 \Omega, \quad \rho_j(x) \xrightarrow{j \rightarrow \infty} 1 \text{ for a.e. } x \in \mathbb{R}^n$$

and $\rho_j f \xrightarrow{j \rightarrow \infty} f$ in $H^1(\Omega)$ for any $f \in H^1(\Omega)$.

Proof. Let $\Gamma_1, \dots, \Gamma_K$ be compact $(n-2)$ -dimensional submanifolds whose union contains $\partial_0 \Omega$ and denote by $d_k(x)$ the distance from $x \in \mathbb{R}^n$ to Γ_k .

Let $f \in H^1(\Omega)$ and $j \in \mathbb{N}$. Denote $h_0 := f$, then for each $k \in \{1, \dots, K\}$ successively use Lemma 13 to choose $j_k \geq j$ such that the function $h_k := (\psi_{j_k} \circ d_k)h_{k-1}$ satisfies

$$\|h_k - h_{k-1}\|_{H^1(\Omega)} < \frac{1}{jK}.$$

By construction one has

$$h_K = \rho_j f, \quad \rho_j := \prod_{k=1}^K (\psi_{j_k} \circ d_k),$$

$$\|\rho_j f - f\|_{H^1(\Omega)} \equiv \|h_K - h_0\|_{H^1(\Omega, \mathbb{C}^N)} \leq \sum_{k=1}^K \|h_k - h_{k-1}\|_{H^1(\Omega, \mathbb{C}^N)} < K \cdot \frac{1}{jK} = \frac{1}{j},$$

hence, $\rho_j f \xrightarrow{j \rightarrow \infty} f$ in $H^1(\Omega)$. The function $\psi_{j_k} \circ d_k$ vanishes in an open neighborhood of Γ_k , so ρ_j vanishes in the union of these neighborhoods (which is also an open neighborhood of $\partial_0 \Omega$). For $j \rightarrow \infty$ one has $j_k \rightarrow \infty$ for each k , hence, $\psi_{j_k} \circ d_k(x) \rightarrow 1$ for $x \notin \Gamma_k$, which implies $\rho_j(x) \xrightarrow{j \rightarrow \infty} 1$ for any $x \notin \bigcup_{k=1}^K \Gamma_k$. Finally, the required bound $0 \leq \rho_j \leq 1$ follows from $0 \leq \psi_j \leq 1$ (Lemma 12). $\square$

Corollary 15. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary and $m \in \mathbb{R}$, then the Dirac operator A_m^Ω is essentially self-adjoint on*

$$\text{dom}_0 A_m^\Omega := \{f \in \text{dom } A_m^\Omega : f = 0 \text{ in an open neighborhood of } \partial_0 \Omega\}.$$

Proof. Let ρ_j be as in Corollary 14 and $f \in \text{dom } A_m^\Omega$. As the boundary condition for A_m^Ω is invariant under multiplication by smooth scalar functions, one has $\rho_j f \in \text{dom}_0 A_m^\Omega$ with $\rho_j f \xrightarrow{j \rightarrow \infty} f$ in $H^1(\Omega, \mathbb{C}^N)$, and the conclusion follows by Lemma 9. $\square$

Lemma 16 (Essential self-adjointness on smooth functions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary, then for any $m \in \mathbb{R}$ the Dirac operator A_m^Ω is essentially self-adjoint on*

$$\text{dom}_\infty A_m^\Omega := \{f \in \text{dom } A_m^\Omega : f \in C^\infty(\overline{\Omega}, \mathbb{C}^N), f = 0 \text{ in an open neighborhood of } \partial_0 \Omega\}.$$

Proof. Let $f \in \text{dom } A_m^\Omega$ and $\varepsilon > 0$. By Corollary 15 one can find a function $h \in \text{dom } A_m^\Omega$ which vanishes in an open neighborhood U of $\partial_0 \Omega$ and satisfies $\|h - f\|_{H^1(\Omega, \mathbb{C}^N)} < \frac{\varepsilon}{3}$. Choose bounded open neighborhoods V_0 and V of $\partial_0 \Omega$ such that $\overline{V_0} \subset V$ and $\overline{V} \subset U$, then pick a function $\chi \in C_c^\infty(\mathbb{R}^n)$ with $0 \leq \chi \leq 1$ such that $\text{supp } \chi \subset V$ and $\chi = 1$ in V_0 . The map

$$H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N) \ni \varphi \mapsto (1 + i\beta\alpha \cdot \nu)(1 - \chi)\varphi \in H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N)$$

is a bounded linear operator (being a multiplication by a smooth matrix function compactly supported on the regular boundary), and we denote its norm by C_0 . Furthermore, it maps the subspace

$$\{\varphi \in H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N) : \text{supp } \varphi \subset \partial_\infty\Omega, \varphi \in C^\infty(\partial_\infty\Omega, \mathbb{C}^N)\}$$

in itself and, in addition, $(1 + i\beta\alpha \cdot \nu)(1 - \chi)\varphi = (1 + i\beta\alpha \cdot \nu)\varphi$ for $\text{supp } \varphi \subset \partial\Omega \setminus V_0$.

Now let Ω_0 be a bounded C^∞ -smooth domain which coincides with Ω outside V_0 (see Lemma 11). For any $\varphi \in H^{\frac{1}{2}}(\partial\Omega_0)$ there is a unique $F \in H^1(\Omega_0)$ with $\Delta F = 0$ in Ω_0 and $F = \varphi$ on $\partial\Omega_0$, and the associated Poisson operator $\mathcal{E}_0 : H^{\frac{1}{2}}(\partial\Omega_0) \ni \varphi \mapsto F \in H^1(\Omega_0)$ is bounded with $\mathcal{E}_0(C^\infty(\partial\Omega_0)) \subset C^\infty(\overline{\Omega_0})$, see [23, Theorem 11.14]. Remark that for any function φ defined on $\partial\Omega$ we can consider $(1 - \chi)\varphi$ as a function defined on $\partial\Omega_0$ and vanishing on $V_0 \cap \partial\Omega_0$. Similarly, for any function v_0 on Ω_0 one can consider $(1 - \chi)v_0$ as a function on Ω which vanishes in $V_0 \cap \Omega$. With these identifications we conclude that

$$\mathcal{E} : H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N) \ni \varphi \mapsto (1 - \chi)(\mathcal{E}_0 \otimes I_N)((1 - \chi)\varphi) \in H^1(\Omega, \mathbb{C}^N)$$

is a bounded operator, and we denote by C_1 its norm. Moreover, it maps $C^\infty(\partial\Omega)$ to $C^\infty(\overline{\Omega})$ by construction, and $\mathcal{E}\varphi = \varphi$ on $\partial\Omega$ for any $\varphi \in C^\infty(\partial\Omega)$ with $\varphi = 0$ on $V \cap \partial\Omega$.

Let $\tilde{h} \in H^1(\mathbb{R}^n, \mathbb{C}^N)$ be an arbitrary continuation of the above h . Consider its convolutions $\tilde{h}_\varepsilon := \tilde{h} * \rho_\varepsilon$ with standard mollifiers ρ_ε , and let h_ε be their restrictions to Ω , then $h_\varepsilon \in C^\infty(\overline{\Omega}, \mathbb{C}^N)$. Having in mind that $h = 0$ in U and the support of h_ε is contained in the ε -neighborhood of the support of h we conclude that we can choose ε sufficiently small such that: the function $g := h_\varepsilon \in C^\infty(\overline{\Omega}, \mathbb{C}^N)$ satisfies $g = 0$ on $V \cap \Omega$ with

$$\|g - h\|_{H^1(\Omega, \mathbb{C}^N)} < \frac{\varepsilon}{3}, \quad \|g - h\|_{H^{\frac{1}{2}}(\partial\Omega)} < \frac{\varepsilon}{3C_0C_1}.$$

Consider the function $v := g - \mathcal{E}(1 + i\beta\alpha \cdot \nu)(g|_{\partial\Omega})$. Due to the above considerations we have $v \in C^\infty(\overline{\Omega}, \mathbb{C}^N)$ with $v = 0$ in V_0 , and $(1 + i\beta\alpha \cdot \nu)(v|_{\partial\Omega}) = 0$, i.e. $v \in \text{dom}_\infty A$. In addition, using $(1 + i\beta\alpha \cdot \nu)(h|_{\partial\Omega}) = 0$ we obtain

$$\begin{aligned} \|v - g\|_{H^1(\Omega, \mathbb{C}^N)} &= \left\| \mathcal{E}(1 + i\beta\alpha \cdot \nu)(g|_{\partial\Omega}) \right\|_{H^1(\Omega, \mathbb{C}^N)} \leq C_1 \left\| (1 + i\beta\alpha \cdot \nu)(g|_{\partial\Omega}) \right\|_{H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N)} \\ &= C_1 \left\| (1 + i\beta\alpha \cdot \nu)(1 - \chi)(g|_{\partial\Omega} - h|_{\partial\Omega}) \right\|_{H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N)} \leq C_1 C_0 \|g - h\|_{H^{\frac{1}{2}}(\partial\Omega, \mathbb{C}^N)} < \frac{\varepsilon}{3}, \end{aligned}$$

therefore,

$$\|v - f\|_{H^1(\Omega, \mathbb{C}^N)} \leq \|v - g\|_{H^1(\Omega, \mathbb{C}^N)} + \|g - h\|_{H^1(\Omega, \mathbb{C}^N)} + \|h - f\|_{H^1(\Omega, \mathbb{C}^N)} < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon,$$

and the claims follows due to the arbitrariness of ε . $\square$

3. PROOF OF THEOREM 3 (QUADRATIC FORM)

As a simple application of the preceding approximations we show that the formulas for the quadratic forms of A_m^2 and $B_{m,M}^2$ previously obtained for smooth Ω are also valid for convex piecewise smooth Ω . The following assertion completes the proof of Theorem 3:

Lemma 17. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary, then for any $f \in \text{dom } A_m^\Omega$ one has*

$$\|A_m^\Omega f\|_{L^2(\Omega, \mathbb{C}^N)}^2 = \int_\Omega (|\nabla f|^2 + m^2|f|^2) dx + \int_{\partial\Omega} \left(m + \frac{H}{2}\right) |f|^2 d\mathcal{H}^{n-1},$$

where H is the mean curvature on $\partial_\infty\Omega$.

Proof. Let $f \in \text{dom}_0 A_m^\Omega$ (see Lemma 16) vanish in an open neighborhood V_0 of $\partial_0 \Omega$. By Lemma 11 one can find a bounded C^∞ -smooth domain Ω_0 coinciding with Ω outside V_0 . Define $f_0 : \Omega_0 \rightarrow \mathbb{C}^N$ by $f_0 := f$ in $\Omega_0 \setminus V_0 \equiv \Omega \setminus V_0$ and $f_0 := 0$ in $\Omega_0 \cap V_0$, then $f_0 \in \text{dom } A_m^{\Omega_0}$ with

$$\begin{aligned} \nabla f_0 &= \nabla f \text{ and } A_m^{\Omega_0} f_0 = A_m^\Omega f \text{ in } \Omega_0 \setminus V_0 \equiv \Omega \setminus V_0, \\ \nabla f_0 &= 0 \text{ and } A_m^{\Omega_0} f_0 = 0 \text{ in } \Omega_0 \cap V_0, \quad \nabla f = 0 \text{ and } A_m^\Omega f = 0 \text{ in } \Omega \cap V_0, \\ f_0 &= f \text{ on } \partial\Omega_0 \setminus V_0 \equiv \partial\Omega \setminus V_0, \quad f_0 = 0 \text{ on } \partial\Omega_0 \cap V, \quad f = 0 \text{ on } \partial\Omega \cap V. \end{aligned}$$

In addition, the mean curvature H_0 on $\partial\Omega_0$ satisfies $H_0 = H$ on $\partial\Omega_0 \setminus V_0 \equiv \partial\Omega \setminus V_0$. Hence,

$$\begin{aligned} \|A_m^\Omega f\|_{L^2(\Omega, \mathbb{C}^N)}^2 &= \|A_m^{\Omega_0} f\|_{L^2(\Omega_0 \setminus V_0, \mathbb{C}^N)}^2 = \|A_m^{\Omega_0} f_0\|_{L^2(\Omega_0 \setminus V_0, \mathbb{C}^N)}^2 = \|A_m^{\Omega_0} f_0\|_{L^2(\Omega_0, \mathbb{C}^N)}^2 \\ (\text{use Lemma 5}) &= \int_{\Omega_0} \left(|\nabla f_0|^2 + m^2 |f_0|^2 \right) dx + \int_{\partial\Omega_0} \left(m + \frac{H_0}{2} \right) |f_0|^2 d\mathcal{H}^{n-1} \\ &= \int_{\Omega_0 \setminus V_0} \left(|\nabla f_0|^2 + m^2 |f_0|^2 \right) dx + \int_{\partial\Omega_0 \setminus V_0} \left(m + \frac{H_0}{2} \right) |f_0|^2 d\mathcal{H}^{n-1} \\ &= \int_{\Omega \setminus V_0} \left(|\nabla f|^2 + m^2 |f|^2 \right) dx + \int_{\partial\Omega \setminus V_0} \left(m + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} \\ &= \int_{\Omega} \left(|\nabla f|^2 + m^2 |f|^2 \right) dx + \int_{\partial\Omega} \left(m + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1}. \end{aligned}$$

Hence the required formula holds for all functions in $\text{dom}_0 A_m^\Omega$. Remark that H can become unbounded near $\partial_0 \Omega$, so some attention is needed when extending to the whole domain.

Let $f \in \text{dom } A_m^\Omega$ and ρ_j be as in Corollary 15, then for $f_j := \rho_j f \in \text{dom}_0 A_m^\Omega$ we have

$$\begin{aligned} \|A_m^\Omega f\|_{L^2(\Omega, \mathbb{C}^N)}^2 &= \lim_{j \rightarrow \infty} \|A_m^\Omega f_j\|_{L^2(\Omega, \mathbb{C}^N)}^2 \\ &= \lim_{j \rightarrow \infty} \left[\int_{\Omega} \left(|\nabla f_j|^2 + m^2 |f_j|^2 \right) dx + \int_{\partial\Omega} \left(m + \frac{H}{2} \right) |f_j|^2 d\mathcal{H}^{n-1} \right]. \end{aligned}$$

By Corollary 15 we have

$$\begin{aligned} \int_{\Omega} \left(|\nabla f_j|^2 + m^2 |f_j|^2 \right) dx &\xrightarrow{j \rightarrow \infty} \int_{\Omega} \left(|\nabla f|^2 + m^2 |f|^2 \right) dx, \\ \int_{\partial\Omega} m |f_j|^2 d\mathcal{H}^{n-1} &\xrightarrow{j \rightarrow \infty} \int_{\partial\Omega} m |f|^2 d\mathcal{H}^{n-1}. \end{aligned}$$

Furthermore, $|f_j| \leq |f|$ with $f_j(x) \xrightarrow{j \rightarrow \infty} f(x)$ for a.e. x , and Fatou's lemma yields

$$\int_{\partial\Omega} \frac{H}{2} |f_j|^2 d\mathcal{H}^{n-1} \xrightarrow{j \rightarrow \infty} \int_{\partial\Omega} \frac{H}{2} |f|^2 d\mathcal{H}^{n-1},$$

which concludes the proof. $\square$

As a preparation for the next section we use similar ideas to show an analogous result for the Dirac operator $B_{m,M}^\Omega$ on $\mathbb{R}^n$ defined in (4).

Lemma 18. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain with piecewise smooth boundary and $m, M \in \mathbb{R}$, then for all $f \in \text{dom } B_{m,M}^\Omega$ it holds that*

$$\begin{aligned} \|B_{m,M}^\Omega f\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 &= \int_{\Omega} \left(|\nabla f|^2 + m^2 |f|^2 \right) dx + \int_{\Omega^c} \left(|\nabla f|^2 + M^2 |f|^2 \right) dx \\ &\quad + (M - m) \int_{\partial\Omega} \left(|\mathcal{P}_- f|^2 - |\mathcal{P}_+ f|^2 \right) d\mathcal{H}^{n-1}, \quad (18) \end{aligned}$$

where $\mathcal{P}_\pm(s) := \frac{1 \mp i\beta\alpha \cdot \nu(s)}{2}$ for $s \in \partial_\infty \Omega$.

Proof. Recall that the formula is already proved for smooth Ω in [30, Lemma 2.3]. Assume first that $f \in H^1(\mathbb{R}^n, \mathbb{C}^N)$ and vanishes in an open neighborhood V_0 of $\partial_0\Omega$. By Lemma 11 one can find a bounded C^∞ -smooth domain Ω_0 which coincides with Ω outside V_0 . Let ν_0 be the outer unit normal on $\partial\Omega_0$ and consider the respective pointwise projectors

$$\mathcal{P}_\pm^0(s) := \frac{1 \mp i\beta\alpha \cdot \nu_0(s)}{2}, \quad s \in \partial\Omega_0,$$

then $\mathcal{P}_\pm^0 f = \mathcal{P}_\pm f$ on $\partial\Omega \setminus V_0$. Due to $B_{m,M}^\Omega f = B_{m,M}^{\Omega_0} f$ one can use the already known formula for the smooth Ω_0 as follows:

$$\begin{aligned} \|B_{m,M}^\Omega f\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 &= \|B_{m,M}^{\Omega_0} f\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 \\ &= \int_{\Omega_0} (|\nabla f|^2 + m^2|f|^2) dx + \int_{\Omega_0^c} (|\nabla f|^2 + M^2|f|^2) dx \\ &\quad + (M - m) \int_{\partial\Omega_0} (|\mathcal{P}_-^0 f|^2 - |\mathcal{P}_+^0 f|^2) d\mathcal{H}^{n-1} \\ &= \int_{\Omega_0 \setminus V_0} (|\nabla f|^2 + m^2|f|^2) dx + \int_{\Omega_0^c \setminus V_0} (|\nabla f|^2 + M^2|f|^2) dx \\ &\quad + (M - m) \int_{\partial\Omega_0 \setminus V_0} (|\mathcal{P}_-^0 f|^2 - |\mathcal{P}_+^0 f|^2) d\mathcal{H}^{n-1} \\ &= \int_{\Omega \setminus V_0} (|\nabla f|^2 + m^2|f|^2) dx + \int_{\Omega^c \setminus V_0} (|\nabla f|^2 + M^2|f|^2) dx \\ &\quad + (M - m) \int_{\partial\Omega \setminus V_0} (|\mathcal{P}_- f|^2 - |\mathcal{P}_+ f|^2) d\mathcal{H}^{n-1} \\ &= \int_{\Omega} (|\nabla f|^2 + m^2|f|^2) dx + \int_{\Omega^c} (|\nabla f|^2 + M^2|f|^2) dx \\ &\quad + (M - m) \int_{\partial\Omega} (|\mathcal{P}_- f|^2 - |\mathcal{P}_+ f|^2) d\mathcal{H}^{n-1}. \end{aligned}$$

Hence, the required formula holds for the functions vanishing near $\partial_0\Omega$. By Lemma 12 the operator $B_{m,M}^\Omega$ is essentially self-adjoint on the set of all such functions, which allows one to extend the formula by density to the whole of $\text{dom } B_{m,M}^\Omega$. $\square$

4. PROOF OF THEOREM 4 (INFINITE MASS LIMIT)

In this section the domain Ω and the mass parameter m in Ω are fixed, and we abbreviate

$$A := A_m^\Omega, \quad B_M := B_{m,M}^\Omega.$$

Recall that $\text{dom}_\infty A$ is defined in Lemma 16. Let us briefly discuss some geometric ingredients appearing in the constructions below. For $s \in \partial_\infty\Omega$ the shape operator $W(s) : T_s\partial_\infty\Omega \rightarrow T_s\partial_\infty\Omega$ is defined by $W(s) := d|_s\nu$. Its eigenvalues $\kappa_1(s), \dots, \kappa_{n-1}(s)$ are the principal curvatures of the boundary at s . One often uses the Jacobian (with I being the identity operator and $t > 0$)

$$J(s, t) := \det(I + tW(s)) \equiv \prod_{j=1}^{n-1} (1 + t\kappa_j(s)) \equiv 1 + \sum_{k=1}^{n-1} H_k(s)t^k, \quad (19)$$

$$\text{with } H_k(s) := \sum_{1 \leq j_1 < \dots < j_k \leq n-1} \kappa_{j_1}(s) \cdot \dots \cdot \kappa_{j_k}(s), \quad k \in \{1, \dots, n-1\}.$$

In particular, $H_1 = \kappa_1 + \dots + \kappa_{n-1} \equiv H$.

4.1. Upper bound. Let $j \in \mathbb{N}$ and denote $E := E_j(A^2)$. Let $\varepsilon > 0$, then by Lemma 16 and min-max principle one can find a j -dimensional subspace $F \subset \text{dom}_\infty A$ such that

$$\max_{f \in F \setminus \{0\}} \frac{\|Af\|_{L^2(\Omega, \mathbb{C}^N)}^2}{\|f\|_{L^2(\Omega, \mathbb{C}^N)}^2} \leq E + \varepsilon. \quad (20)$$

Due to convexity the map $S : \Omega^\mathbb{G} \rightarrow \partial\Omega$ given by the rule $|x - S(x)| \leq |x - y|$ for any $y \in \partial\Omega$ is a well-defined Lipschitz function. For any $f \in F$ denote by $\tilde{f}$ its extension to $\mathbb{R}^n$ by

$$\tilde{f}(x) := f(S(x))e^{-Md(x)}, \quad d(x) := \text{dist}(x, \partial\Omega), \quad x \in \Omega^\mathbb{G},$$

and consider the j -dimensional subspace $\tilde{F} := \{\tilde{f} : f \in F\} \subset H^1(\mathbb{R}^n, \mathbb{C}^N)$. Our goal is to show that for some $c > 0$ and all sufficiently large $M > 0$ one has

$$\max_{\tilde{f} \in \tilde{F} \setminus \{0\}} \frac{\|B_M \tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2}{\|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2} \leq E + \varepsilon + \frac{c}{M}, \quad (21)$$

then the min-max principle implies first $\limsup_{M \rightarrow +\infty} E_j(B_M^2) \leq E + \varepsilon$, and due to the arbitrariness of ε one obtains the upper bound

$$\limsup_{M \rightarrow +\infty} E_j(B_M^2) \leq E \equiv E_j(A^2). \quad (22)$$

Now let us show (21). For any $\tilde{f} \in \tilde{F}$ one has $\mathcal{P}_+ \tilde{f} = \tilde{f}$ and $\mathcal{P}_- \tilde{f} = 0$, hence, with $f := \tilde{f}|_\Omega \in F \subset \text{dom}_\infty A$ one can rewrite the expression of Lemma 18 in the form

$$\begin{aligned} \|B_M \tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 &\equiv \int_\Omega (|\nabla f|^2 + m^2|f|^2) dx + \int_{\partial\Omega} \left(m + \frac{H_1}{2}\right) |f|^2 d\mathcal{H}^{n-1} \\ &\quad + \int_{\Omega^\mathbb{G}} (|\nabla \tilde{f}|^2 + M^2|u|^2) dx - \int_{\partial\Omega} \left(M + \frac{H_1}{2}\right) |\tilde{f}|^2 d\mathcal{H}^{n-1} \\ &\equiv \|Af\|_{L^2(\Omega, \mathbb{C}^N)}^2 + R_M(\tilde{f}), \\ R_M(\tilde{f}) &:= \int_{\Omega^\mathbb{G}} (|\nabla \tilde{f}|^2 + M^2|\tilde{f}|^2) dx - \int_{\partial\Omega} \left(M + \frac{H_1}{2}\right) |\tilde{f}|^2 d\mathcal{H}^{n-1}. \end{aligned} \quad (23)$$

Let us derive an upper bound for $R_M(\tilde{f})$ using local coordinates. Remark first that one can find a compact subset $K \subset \partial_\infty \Omega$ with $\text{supp } \tilde{f}|_{\partial\Omega} \subset K$ for all $\tilde{f} \in \tilde{F}$, and then $\tilde{f}(x) = 0$ for all $x \in \Omega^\mathbb{G}$ with $S(x) \notin K$. We further choose open subsets $V_1, \dots, V_P \subset \partial_\infty \Omega$ such that $K \subset V_1 \cup \dots \cup V_P$ and each V_p is covered by a local chart $\varphi_p : \mathbb{R}^{n-1} \supset U_p \rightarrow V_p$, and we pick $\chi_p \in C_c^\infty(V_p)$ such that $\chi_1 + \dots + \chi_P = 1$ on K . Then $\chi_1 \circ S + \dots + \chi_P \circ S = 1$ on $\text{supp } \tilde{f} \cap \Omega^\mathbb{G}$ for all $\tilde{f} \in \tilde{F}$. Furthermore, the maps

$$\Phi_p : U_p \times (0, \infty) \ni (s, t) \mapsto \varphi_p(s) + t\nu(\varphi_p(s)) \in \Omega^\mathbb{G}$$

are diffeomorphisms satisfying the identities $d(\Phi_p(s, t)) = t$ and $S(\Phi_p(s, t)) = \varphi_p(s)$.

Let $\tilde{f} \in \tilde{F}$ and denote $f_p := \tilde{f} \circ \Phi_p$, then $f_p(s, t) = f(\varphi_p(s))e^{-Mt}$ for all (s, t) , and the partition of unity $\chi_p \circ S$ and a standard change of variables yield:

$$\begin{aligned} R_M(\tilde{f}) &= \sum_{p=1}^P \left[\int_{\Omega^\mathbb{G}} (\chi_p \circ S) (|\nabla \tilde{f}|^2 + M^2|\tilde{f}|^2) dx - \int_{\partial\Omega} \chi_p \left(M + \frac{H_1}{2}\right) |\tilde{f}|^2 d\mathcal{H}^{n-1} \right] \\ &= \sum_{p=1}^P \int_{U_p} \chi_p(\varphi_p(s)) \left[\int_0^\infty \left(\langle \nabla f_p(s, t), G_p(s, t)^{-1} \nabla f_p(s, t) \rangle + M^2 |f_p(s, t)|^2 \right) J_p(s, t) dt \right] \end{aligned}$$

$$- \left(M + \frac{H_1(\varphi_p(s))}{2} \right) |f_p(s, 0)|^2 \Big] \sqrt{\det g_p(s)} \, ds,$$

with matrices

$$G_p(s, t) := \begin{pmatrix} \left(\left\langle (1 + tW_p(s)) \partial_j \varphi(s), (1 + tW_p(s)) \partial_k \varphi(s) \right\rangle \right)_{j,k=1}^{n-1} & 0 \\ 0 & 1 \end{pmatrix},$$

$$g_p(s) := \left(\left\langle \partial_j \varphi(s), \partial_k \varphi(s) \right\rangle \right)_{j,k=1}^{n-1}, \quad W_p := W \circ \varphi_p.$$

and Jacobians $J_p(s, t) := J(\varphi_p(s), t)$, see (19). As $W_p(s)$ is self-adjoint and its eigenvalues (the principal curvatures) are non-negative due to convexity, the min-max principle implies $G_p(s, t) \geq G_p(s, 0) \equiv g_p(s) \oplus 1$ resulting in

$$G_p(s, t)^{-1} \leq G_p(s, 0)^{-1} \equiv \begin{pmatrix} g_p(s)^{-1} & 0 \\ 0 & 1 \end{pmatrix}.$$

Hence,

$$\begin{aligned} R_M(\tilde{f}) &\leq \sum_{p=1}^P \int_{U_p} \chi_p(\varphi_p(s)) \left[\int_0^\infty \left(\left\langle \nabla_s f_p(s, t), g_p^{-1}(s) \nabla_s f_p(s, t) \right\rangle + |\partial_t f_p(s, t)|^2 \right. \right. \\ &\quad \left. \left. + M^2 |f_p(s, t)|^2 \right) J_p(s, t) \, dt - \left(M + \frac{H_1(\varphi_p(s))}{2} \right) |f_p(s, 0)|^2 \right] \sqrt{\det g_p(s)} \, ds \\ &= \sum_{p=1}^P \int_{U_p} \chi_p(\varphi_p(s)) \left[\int_0^\infty \left(\left\langle \nabla_s f(\varphi_p(s)), g_p^{-1}(s) \nabla_s f(\varphi_p(s)) \right\rangle e^{-2Mt} \right. \right. \\ &\quad \left. \left. + 2M^2 |f(\varphi_p(s))|^2 e^{-2Mt} \right) J_p(s, t) \, dt - \left(M + \frac{H_1(\varphi_p(s))}{2} \right) |f(\varphi_p(s))|^2 \right] \\ &\quad \times \sqrt{\det g_p(s)} \, ds. \end{aligned}$$

A simple direct computation shows that

$$\int_0^\infty e^{-2Mt} \, dt = \frac{1}{2M}, \quad \int_0^\infty J_p(s, t) e^{-2Mt} \, dt = \frac{1}{2M} + \sum_{k=1}^{n-1} \frac{k! H_k(\varphi_p(s))}{(2M)^{k+1}},$$

and there is a $C_p > 0$ such that for all s with $\varphi_p(s) \in \text{supp } \chi_p$ and sufficiently large M one has

$$\int_0^\infty J_p(s, t) e^{-2Mt} \, dt \leq \frac{1}{2M^2} \left(M + \frac{H_1(\varphi_p(s))}{2} \right) + \frac{C_p}{M^3},$$

which results in

$$\begin{aligned} R_M(\tilde{f}) &\leq \sum_{p=1}^P \int_{U_p} \chi_p(\varphi_p(s)) \left[\frac{1}{2M} \left\langle \nabla_s f(\varphi_p(s)), g_p^{-1}(s) \nabla_s f(\varphi_p(s)) \right\rangle \right. \\ &\quad \left. + \frac{2C_p}{M} |f(\varphi_p(s))|^2 \right] \sqrt{\det g_p(s)} \, ds \leq \frac{C}{M} \|\tilde{f}\|_{H^1(\partial\Omega, \mathbb{C}^N)}^2 \end{aligned}$$

for some constant $C > 0$ independent of $\tilde{f}$ and M . In addition, one can find another constant $C_1 > 0$ with $\|\tilde{f}\|_{H^1(\partial\Omega, \mathbb{C}^N)}^2 \leq C_1 \|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2$ for all $\tilde{f} \in \tilde{F}$, as $\tilde{F}$ is finite-dimensional. Hence,

$$R_M(\tilde{f}) \leq \frac{CC_1}{M} \|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 \quad \text{for all } \tilde{f} \in \tilde{F}.$$

By inserting this estimate and (20) into the representation (23) we obtain for all $\tilde{f} \in \tilde{F} \setminus \{0\}$:

$$\frac{\|B_M \tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2}{\|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2} \leq \frac{(E + \varepsilon)\|f\|_{L^2(\Omega, \mathbb{C}^N)}^2 + \frac{CC_1}{M}\|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2}{\|\tilde{f}\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2} \leq E + \varepsilon + \frac{CC_1}{M},$$

which shows the sought estimate (21) with $c := CC_1$.

We make a side remark that the upper bound did not require the boundedness of H .

4.2. Lower bound. We start with a preliminary estimate:

Lemma 19. *Let the mean curvature H of $\partial_\infty \Omega$ be bounded, then for some $c > 0$ one has*

$$R_\gamma(f) := \int_{\Omega^c} (|\nabla f|^2 + \gamma^2 |f|^2) dx - \int_{\partial\Omega} \left(\gamma + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} \geq -c \int_{\Omega^c} |f|^2 dx$$

for all $f \in H^1(\Omega^c)$ and $\gamma > 0$.

Proof. We adapt some constructions from [32, Thm. 2]. Denote $\Pi := \partial_\infty \Omega \times (0, \infty)$ and remark that the map $\Phi : \Pi \ni (s, t) \mapsto s + t\nu(s)$ is a diffeomorphism between Π and $\Phi(\Pi) \subset \Omega^c$. For any $f \in C_c^\infty(\mathbb{R}^n)$ one has, using the standard change of variables,

$$\begin{aligned} R_\gamma(f) &\geq \int_{\Phi(\Pi)} (|\nabla f|^2 + \gamma^2 |f|^2) dx - \int_{\partial_\infty \Omega} \left(\gamma + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} \\ &= \int_{\partial_\infty \Omega} \int_0^\infty \left(|(\nabla f)(\Phi(s, t))|^2 + \gamma^2 |f(\Phi(s, t))|^2 \right) J(s, t) dt d\mathcal{H}^{n-1}(s) \\ &\quad - \gamma \int_{\partial_\infty \Omega} \left(\gamma + \frac{H_1(s)}{2} \right) |f(\Phi(s, 0))|^2 d\mathcal{H}^{n-1}(s) \end{aligned}$$

with J given by (19). Furthermore,

$$|(\nabla f)(\Phi(s, t))|^2 \geq \left| \left\langle \nu(s), (\nabla f)(\Phi(s, t)) \right\rangle \right|^2 = \left| \frac{\partial}{\partial t} f(\Phi(s, t)) \right|^2, \quad (24)$$

hence,

$$\begin{aligned} R_\gamma(f) &\geq R'_\gamma(f) := \int_{\partial_\infty \Omega} \left[\int_0^\infty \left(\left| \frac{\partial}{\partial t} g(s, t) \right|^2 + \gamma^2 |g(s, t)|^2 \right) J(s, t) dt \right. \\ &\quad \left. - \left(\gamma + \frac{H_1}{2} \right) |g(s, 0)|^2 \right] d\mathcal{H}^{n-1}(s) \quad \text{with } g := f \circ \Phi. \end{aligned} \quad (25)$$

We further consider the function $h := \sqrt{J}g$, then

$$\begin{aligned} |\partial_t g|^2 &= \left| \partial_t \frac{h}{\sqrt{J}} \right|^2 = \left| \frac{\partial_t h}{\sqrt{J}} - \frac{h \partial_t J}{2J\sqrt{J}} \right|^2 = \frac{1}{J} \left| \partial_t h - \frac{\partial_t J}{2J} h \right|^2 \\ &= \frac{1}{J} \left(|\partial_t h|^2 - \frac{\partial_t J}{2J} \partial_t |h|^2 + \left(\frac{\partial_t J}{2J} \right)^2 |h|^2 \right), \end{aligned}$$

therefore,

$$\begin{aligned} R'_\gamma(f) &= \int_{\partial_\infty \Omega} \left[\int_0^\infty \left(|\partial_t h|^2 - \frac{\partial_t J}{2J} \partial_t |h|^2 + \left(\frac{\partial_t J}{2J} \right)^2 |h|^2 + \gamma^2 |h|^2 \right) dt \right. \\ &\quad \left. - \left(\gamma + \frac{H_1(s)}{2} \right) |h(s, 0)|^2 \right] d\mathcal{H}^{n-1}(s). \end{aligned}$$

Remark that

$$\begin{aligned} \frac{\partial_t J(s, t)}{2J(s, t)} &= \frac{1}{2} \partial_t \log J(s, t) = \frac{1}{2} \partial_t \sum_{j=1}^{n-1} \log(1 + t\kappa_j(s)) = \frac{1}{2} \sum_{j=1}^{n-1} \frac{\kappa_j(s)}{1 + t\kappa_j(s)}, \\ \left. \frac{\partial_t J(s, t)}{2J(s, t)} \right|_{t=0} &= \frac{1}{2} H_1(s), \end{aligned} \quad (26)$$

and the integration by parts yields

$$\begin{aligned} \int_0^\infty \left(-\frac{\partial_t J(s, t)}{2J(s, t)} \partial_t |h(s, t)|^2 \right) dt &= \left[-\frac{\partial_t J(s, t)}{2J(s, t)} |h(s, t)|^2 \right]_{t=0}^{t=\infty} + \int_0^\infty \partial_t \left(\frac{\partial_t J(s, t)}{2J(s, t)} \right) |h(s, t)|^2 dt \\ &= \frac{H_1(s)}{2} |h(s, 0)|^2 + \int_0^\infty \partial_t \left(\frac{\partial_t J(s, t)}{2J(s, t)} \right) |h(s, t)|^2 dt, \end{aligned}$$

and the substitution into the preceding expression for $R'_\gamma(f)$ gives

$$\begin{aligned} R'_\gamma(f) &= \int_{\partial_\infty \Omega} \left[\int_0^\infty \left(|\partial_t h(s, t)|^2 + \gamma^2 |h(s, t)|^2 \right) dt - \gamma |h(s, 0)|^2 \right] d\mathcal{H}^{n-1}(s) \\ &\quad + \int_{\partial_\infty \Omega} \int_0^\infty \left(\partial_t \left(\frac{\partial_t J}{2J} \right) + \left(\frac{\partial_t J}{2J} \right)^2 \right) |h|^2 dt d\mathcal{H}^{n-1}. \end{aligned} \quad (27)$$

Using

$$\begin{aligned} |h(s, 0)|^2 &= -\int_0^\infty \partial_t |h(s, t)|^2 dt = -\int_0^\infty 2\Re \left(\overline{\partial_t h(s, t)} h(s, t) \right) dt \\ &\leq \int_0^\infty 2|\partial_t h(s, t)| \cdot |h(s, t)| dt \leq \int_0^\infty \left(\frac{1}{\gamma} |\partial_t h(s, t)|^2 + \gamma |h(s, t)|^2 \right) dt, \end{aligned}$$

we show that the first integral in (27) is always non-negative. To estimate the second integral in (27) we use (26):

$$\begin{aligned} \partial_t \left(\frac{\partial_t J(s, t)}{2J(s, t)} \right) &= -\frac{1}{2} \sum_{j=1}^{n-1} \frac{\kappa_j(s)^2}{(1 + t\kappa_j(s))^2} \geq -\frac{n-1}{2} H(s)^2, \\ \partial_t \left(\frac{\partial_t J}{2J} \right) + \left(\frac{\partial_t J}{2J} \right)^2 &\geq \partial_t \left(\frac{\partial_t J}{2J} \right) \geq -c, \quad c := \frac{n-1}{2} \|H\|_\infty^2. \end{aligned}$$

The substitution into (27) gives

$$\begin{aligned} R'_\gamma(f) &\geq -c \int_{\partial_\infty \Omega} \int_0^\infty |h|^2 dt d\mathcal{H}^{n-1}(s) = -c \int_{\partial_\infty \Omega} \int_0^\infty \left| f(\Phi(s, t)) \right|^2 J(s, t) dt d\mathcal{H}^{n-1}(s) \\ &= -c \int_{\Phi(\Pi)} |f|^2 dx \geq -c \int_{\Omega^{\mathbb{C}}} |f|^2 dx, \end{aligned}$$

and using (25) we arrive at

$$R_\gamma(f) \geq -c \int_{\Omega^{\mathbb{C}}} |f|^2 dx \text{ for all } f \in C_c^\infty(\mathbb{R}^n),$$

which extends by density to all $f \in H^1(\Omega^{\mathbb{C}})$. $\square$

With the lower bound of Lemma 19 at hand, the remaining analysis is very close to the constructions in [30, Sec. 5.3]. We use the formula (18) for $\|B_M f\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2$ to obtain

$$\begin{aligned} \|B_M f\|_{L^2(\mathbb{R}^n, \mathbb{C}^N)}^2 &= \int_{\Omega} (|\nabla f|^2 + m^2 |f|^2) dx \\ &\quad + \int_{\partial\Omega} \left(m - \frac{1}{\sqrt{M}} + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} + 2(M - m) \int_{\partial\Omega} |\mathcal{P}_- f|^2 d\mathcal{H}^{n-1} \end{aligned}$$

$$\begin{aligned}
& + \left[\int_{\Omega^c} (|\nabla f|^2 + M^2|f|^2) dx - \int_{\partial\Omega} \left(M - \frac{1}{\sqrt{M}} + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} \right] \\
& \geq \int_{\Omega} (|\nabla f|^2 + m^2|f|^2) dx + \int_{\partial\Omega} \left(m - \frac{1}{\sqrt{M}} + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} \\
& \quad + 2(M - m) \int_{\partial\Omega} |\mathcal{P}_- f|^2 d\mathcal{H}^{n-1} + C_M \int_{\Omega^c} |f|^2 dx, \quad C_M := 2\sqrt{M} - \frac{1}{M} - c,
\end{aligned}$$

where the term in the square brackets was estimated from below by Lemma 19. Using the Neumann decoupling of Ω and Ω^c along $\partial\Omega$ and the min-max principle we conclude that for any fixed $j \in \mathbb{N}$ and all $M > 0$ one has

$$E_j(B_M^2) \geq E_j(K_M \oplus C_M I^c), \quad (28)$$

where K_M is the self-adjoint operator with compact resolvent in $L^2(\Omega, \mathbb{C}^N)$ whose sesquilinear form k_M is defined on $\text{dom } k_M := H^1(\Omega, \mathbb{C}^N)$ by

$$\begin{aligned}
k_M(f, f) &= \int_{\Omega} (|\nabla f|^2 + m^2|f|^2) dx \\
&\quad + \int_{\partial\Omega} \left(m - \frac{1}{\sqrt{M}} + \frac{H}{2} \right) |f|^2 d\mathcal{H}^{n-1} + 2(M - m) \int_{\partial\Omega} |\mathcal{P}_- f|^2 d\mathcal{H}^{n-1}
\end{aligned}$$

and I^c is the identity operator in $L^2(\Omega^c, \mathbb{C}^N)$. Due to the upper bound (22) the eigenvalue $E_j(B_M^2)$ remains bounded for $M \rightarrow +\infty$, while $C_M \xrightarrow{M \rightarrow +\infty} +\infty$, and (28) yields

$$\liminf_{M \rightarrow +\infty} E_j(B_M^2) \geq \liminf_{M \rightarrow +\infty} E_j(K_M). \quad (29)$$

For each $f \in H^1(\Omega, \mathbb{C}^N)$ the function $M \rightarrow k_M(f, f)$ is monotonically increasing, with

$$\left\{ f \in \bigcap_{M>0} \text{dom } k_M : \sup_{M>0} k_M(f, f) < \infty \right\} = \left\{ f \in H^1(\Omega, \mathbb{C}^N) : \mathcal{P}_- f = 0 \text{ on } \partial\Omega \right\} = \text{dom } A,$$

and for any $f \in \text{dom } A$ one has $\lim_{M \rightarrow +\infty} k_M(f, f) = \|Af\|_{L^2(\Omega, \mathbb{C}^N)}^2$. Hence, the monotone convergence principle implies $\lim_{M \rightarrow +\infty} E_j(K_M) = E_j(A^2)$, see [35, 39] for the general theory or [30, Prop. 7] for the most adapted formulation. The substitution into (29) gives the required lower bound $\liminf_{M \rightarrow +\infty} E_j(B_M^2) \geq E_j(A^2)$.

Remark 20. While the boundedness of H is explicitly used in Lemma 19, we believe that this assumption is technical and can probably be avoided by applying more advanced proof methods. It is implicitly supported by the fact that the boundary integral containing H in the expression for $\|A_m^\Omega f\|^2$ in Theorem 3 turns out to be always finite, while H can be unbounded. The proof of Lemma 19 does not exploit the tangential derivatives, as they are dropped in the step (24). One may expect that these tangential derivatives can actually compensate the unbounded curvature terms (using the idea that a Schrödinger operator with an unbounded negative potential may still remain lower semibounded). Nevertheless such an improvement would require a very advanced analysis (like a control of unbounded curvatures near the singular boundary), which goes significantly beyond the scope of the present work.

APPENDIX A. DIRAC MATRICES

The most frequently used choices for the Dirac matrices α_k and β entering (1) are based on the Pauli matrices

$$\sigma_1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

For $n = 2$ one has $N = 2$, and the standard choice is $\alpha_k := \sigma_k$ for $k \in \{1, 2\}$ and $\beta := \sigma_3$. For $n = 3$ one has $N = 4$, and one often uses

$$\alpha_k := \begin{pmatrix} 0 & \sigma_k \\ \sigma_k & 0 \end{pmatrix}, \quad k \in \{1, 2, 3\}, \quad \beta := \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix},$$

where I_2 is the 2×2 identity matrix. For higher dimensions one can use various iterative procedures: see e.g. [19, Chapter 15] or [40, Appendix E] for a general theory, or [29, Sec. 2.1] for a more specific description. We remark that the choice of α_k and β is unique up to similarity transforms and sign changes, as described in the aforementioned references, and all associated Dirac operators are unitarily equivalent to each other.

ACKNOWLEDGMENTS

The work was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), project 491606144.

REFERENCES

- [1] P. R. S. Antunes, F. Bento, D. Krejčířík: *Numerical optimisation of Dirac eigenvalues*. J. Phys. A **57** (2024) 475203.
- [2] N. Arrizabalaga, L. Le Treust, A. Mas, N. Raymond: *The MIT bag model as an infinite mass limit*. J. École Polytech. Math. **6** (2019) 329–365.
- [3] N. Arrizabalaga, L. Le Treust, N. Raymond: *Extension operator for the MIT bag model*. Ann. Fac. Sci. Toulouse Math. **29** (2020) 135–147.
- [4] J.-M. Barbaroux, H. D. Cornean, L. Le Treust, E. Stockmeyer: *Resolvent convergence to Dirac operators on planar domains*. Ann. Henri Poincaré **20** (2019) 1877–1891.
- [5] C. Bär, W. Ballmann: *Guide to boundary value problems for Dirac-type operators*. W. Ballmann et al. (Eds.): *Arbeitstagung Bonn 2013*. Progress in Mathematics, Vol. 319, Birkhäuser, 2016, pp. 43–80.
- [6] C. Bär, W. Ballmann: *Boundary value problems for elliptic differential operators of first order*. H.-D. Cao, S.-T. Yau (Eds.): *Algebra and Geometry: In Memory of C. C. Hsiung* (Surveys in Differential Geometry, vol. 17, Intl. Press, 2012), pp. 1–78.
- [7] J. Behrndt, M. Holzmam, A. Mas: *Self-adjoint Dirac operators on domains in $\mathbb{R}^3$* . Ann. Henri Poincaré **21** (2020) 2681–2735.
- [8] J. Behrndt, M. Holzmam, C. Stelzer-Landauer, G. Stenzel: *Boundary triples and Weyl functions for Dirac operators with singular interactions*. Rev. Math. Phys. **36** (2024) 2350036.
- [9] R. D. Benguria, S. Fournais, E. Stockmeyer, H. Van Den Bosch: *Self-adjointness of two-dimensional Dirac operators on domains*. Ann. Henri Poincaré **18** (2017) 1371–1383.
- [10] B. Benhellal: *Spectral analysis of Dirac operators with delta interactions supported on the boundaries of rough domains*. J. Math. Phys. **63** (2022) 011507.
- [11] B. Benhellal, V. Bruneau, M. Zreik: *A Poincaré–Steklov map for the MIT bag model*. Anal. PDE **17** (2024) 2923–2970.
- [12] M. V. Berry, R. J. Mondragon: *Neutrino billiards: time-reversal symmetry-breaking without magnetic fields*. Proc. Roy. Soc. London Ser. A **412** (1987) 53–74.
- [13] P. Bogolioubov: *Sur un modèle à quarks quasi-indépendants*. Ann. IHP A **8** (1968) 163–189.
- [14] B. Booß-Bavnbek, K. P. Wojciechowski: *Elliptic boundary problems for Dirac operators*. Springer, 1993.
- [15] J. Brüning, M. Lesch: *On boundary value problems for Dirac type operators: I. Regularity and self-adjointness*. J. Funct. Anal. **185** (2001) 1–62.
- [16] B. Cassano, V. Lotoreichik: *Self-adjointness for the MIT bag model on an unbounded cone*. Math. Nachr. **297** (2024) 1006–1041.
- [17] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, V. F. Weisskopf: *New extended model of hadrons*. Phys. Rev. D **9** (1974) 3471–3495.
- [18] M. Costabel, M. Dauge, S. Nicaise: *Singularities of Maxwell interface problems*. Math. Model. Num. Anal. **33** (1999) 627–649.
- [19] J. Dereziński, C. Gérard: *Quantization and quantum fields*. Cambridge Univ. Press, 2013.

- [20] B. Flamencourt: *A generalized MIT bag operator on spin manifolds in the non-relativistic limit*. J. Geom. Phys. **178** (2025) 104534.
- [21] P. Grisvard: *Elliptic problems in non-smooth domains*. Pitman Advanced Pub. Program, Boston, 1985.
- [22] N. Große, A. Uribe, H. Van Den Bosch: *Local boundary conditions for Dirac-type operators*. Preprint arXiv:2412.17396.
- [23] G. Grubb: *Distributions and operators*. Springer, 2009.
- [24] K. Johnson: *The MIT bag model*. Acta Phys. Pol. B **6** (1975) 865–892.
- [25] P. D. Lamberti, L. Provenzano: *H^2 -regularity of Steklov eigenfunctions on convex domains via Rellich-Pohozaev identity*. Preprint arXiv:2503.19852.
- [26] P. D. Lamberti, L. Provenzano: *H^2 -regularity on convex domains for Robin eigenfunctions with parameter of arbitrary sign*. Preprint arXiv:2507.16726.
- [27] J. M. Lee: *Introduction to smooth manifolds*. Springer, 2002.
- [28] L. Le Treust, T. Ourmières-Bonafos: *Self-adjointness of Dirac operators with infinite mass boundary conditions in sectors*. Ann. Henri Poincaré **19** (2018) 1465–1487.
- [29] V. Lotoreichik, T. Ourmières-Bonafos: *Spectral asymptotics of the Dirac operator in a thin shell*. J. Math. Anal. Appl. **546** (2025) 129190.
- [30] A. Moroianu, T. Ourmières-Bonafos, K. Pankrashkin: *Dirac operators on hypersurfaces as large mass limits*. Comm. Math. Phys. **374** (2020) 1963–2013.
- [31] T. Ourmières-Bonafos, L. Vega: *A strategy for self-adjointness of Dirac operators: Applications to the MIT bag model and delta-shell interactions*. Publ. Mat. **62** (2018) 397–437.
- [32] K. Pankrashkin: *On the discrete spectrum of Robin Laplacians in conical domains*. Math. Model. Nat. Phenom. **11**:2 (2016) 100–110.
- [33] F. Pizzichillo, H. Van Den Bosch: *Self-adjointness of two dimensional Dirac operators on corner domains*. J. Spectr. Theory **11** (2021) 1043–1079.
- [34] R. Schneider: *Convex bodies: The Brunn–Minkowski theory. Second edition*. Cambridge University Press, 2013.
- [35] B. Simon: *A canonical decomposition for quadratic forms with applications to monotone convergence theorems*. J. Funct. Anal. **28** (1978) 377–385.
- [36] E. Stockmayer, S. Vugalter: *Infinite mass boundary conditions for Dirac operators*. J. Spect. Theory **9** (2019) 569–600.
- [37] B. Thaller: *The Dirac equation*. Springer, 1992.
- [38] T. Vu: *On the regularity of Dirac eigenfunctions*. PDE Appl. **6** (2025) 31.
- [39] J. Weidmann: *Stetige Abhängigkeit der Eigenwerte und Eigenfunktionen elliptischer Differentialoperatoren vom Gebiet*. Math. Scand. **54** (1984) 51–69.
- [40] B. de Wit, J. Smith: *Field theory in particle physics. Volume 1*. North-Holland, Amsterdam etc., 1986.

CARL VON OSSIEZKY UNIVERSITÄT OLDENBURG, FAKULTÄT V – MATHEMATIK UND NATURWISSENSCHAFTEN, INSTITUT FÜR MATHEMATIK, AMMERLÄNDER HEERSTR. 114–118, 26129 OLDENBURG, GERMANY

Email address: `konstantin.pankrashkin@uol.de`

URL: `http://uol.de/pankrashkin`